



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A61K 7/48	A1	(11) International Publication Number: WO 00/67722
(21) International Application Number: PCT/US00/11203 (22) International Filing Date: 26 April 2000 (26.04.00) (30) Priority Data: 09/306,314 6 May 1999 (06.05.99) US (71) Applicant: COLOR ACCESS, INC. [US/US]; 7 Corporate Center Drive, Melville, NY 11747 (US). (72) Inventors: MAMMONE, Thomas; 4 Spencer Street, Farmingdale, NY 11735 (US). GAN, David, C.; 50 Chichester Street, Huntington Station, NY 11746 (US). (74) Agent: TSVEDOS, Estelle, J.; Kenyon & Kenyon, One Broadway, New York, NY 10004 (US).		(81) Designated States: AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.
(54) Title: METHODS OF EXFOLIATING USING N-ACETYL GLUCOSAMINE		
(57) Abstract		
The present invention relates to a method of exfoliating and moisturizing the skin comprising the steps of topically applying a cosmetic or pharmaceutical composition. The composition comprises an amino sugar, N-acetyl-D-glucosamine and related compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS OF EXFOLIATION USING N-ACETYL GLUCOSAMINE

Field of the Invention

- 5 The present invention relates to methods of substantially exfoliating the skin. In particular, the invention relates to topically applied compositions containing an amino sugar in the form of N-acetyl glucosamine.

Background of the Invention

- 10 This invention relates to methods of using N-acetyl glucosamine to exfoliate the skin. The cells of the outermost layer of the stratum corneum are constantly shed naturally, by the normal process of desquamation, as minute particles. When fully keratinised tissue loses its cellular structure and reaches the surface of the stratum corneum, it breaks up into microscopic squames and sheds off the surface of the skin. Microscopic squames at the skin surface are commonly
15 referred to as dead skin cells and make up a dead skin layer on the skin surface. The process of desquamation has been estimated to cause a loss of tissue in an amount of up to 14 grams per day. This loss is constantly replenished with cells from lower layers of the epidermis. Thus, the layers of the epidermis are composed of cells moving up towards the surface in successive stages of differentiation until death when they are finally sloughed off and lost to the environment.
20 Desquamation is one of the processes by which skin maintains its health and vitality as nutrients and moisture are continuously replaced on the surface of the skin when dead skin cells are removed. Normally, the desquamation process takes about 14 days (i.e., the corneocyte takes 14 days to reach the outermost layer of the stratum corneum to be shed). When desquamation does not take place regularly, the surface of the skin tends to become rougher and more wrinkles and
25 other undesirable effects appear on the surface of the skin. However, in addition to or as an alternative to the natural desquamation process, exfoliation is often used to rejuvenate and enhance the health of the skin.

- Exfoliation is a technique whereby dead skin cells are removed or sloughed from the skin surface to promote a healthier and more youthful appearance to the skin. Several compounds are
30 known to be useful as exfoliants such as for example, alpha hydroxy acid ("AHA"), beta hydroxy acid ("BHA"), retinoic acid ("retin A"), and enzymes. An exfoliant such as AHA breaks the bond holding individual squames together and allows them to detach and shed.

Producing a variety of alternatives for exfoliation is desirable because skin types vary among consumers and therefore, having a variety of exfoliants available to meet various individual skincare needs of consumers is beneficial. Thus, there is a continued effort to find additional alternative ways of aiding the sloughing ability of the skin and promoting its health for various types of skin. This is, therefore, an object of the present invention.

Summary of the Invention

The present invention relates to a method for exfoliating the skin by topically applying pharmaceutical or cosmetic compositions containing an exfoliating-effective amount of N-acetyl-D-glucosamine, N-acetylgalactosamine, or a combination thereof. This method provides a natural alternative to the exfoliation process that is as effective as other typical exfoliation techniques.

Detailed Description of the Invention

It has now been unexpectedly discovered that topically applied N-acetyl-D-glucosamine or N-acetylgalactosamine containing compositions exfoliate the skin. Combinations of N-acetyl-D-glucosamine and N-acetylgalactosamine can also be used. These compositions which exfoliate the skin can also include chitin to enhance the process of exfoliation.

It is known that amino sugars are capable of modulating the adhesion of keratinocytes, *in vitro*. It is disclosed by Brysk, M. in "Glycoproteins modulate adhesion in terminally differentiated keratinocytes," that N-acetylglucosamine, N-acetylneuraminic acid and N-acetylgalactosamine are involved in the dissociation of aggregates of glycoproteins that bind corneocytes together. Specifically, amino sugars are known to inhibit the reaggregation of corneocytes which have been dissociated into single squames by homogenization in ether. Further, amino sugars in the form of N-acetylglucosamine have been used as an oral nutrient supplement in conjunction with other topical cosmetic products, such as for example, products offered by Life-Force, Inc. which include N-acetyl-D-glucosamine in the form of a pill taken as a nutritional supplement. However, it is not suggested in the prior art to formulate a pharmaceutical or cosmetic composition for topical application to the skin using N-acetylglucosamine as an exfoliant. In fact, in WO97/12597 it is disclosed that when studied to determine its efficacy as a topical desquamating agent, N-acetyl-D-glucosamine was not found to be effective. The exfoliant in WO97/12597 is a compound comprising a chain of carbohydrates linked by a linking moiety to an alkyl or alkenyl chain. Further, WO97/12597, incorporated herein by reference, only includes N-acetyl-D-glucosamine as one of the units forming the

carbohydrate portion which is linked by the linking moiety to the alkyl or alkenyl chain for desquamation of the skin. It is not disclosed in WO97/12597 that N-acetyl-D-glucosamine alone can exfoliate the skin.

Compositions for topical application containing N-acetyl-D-glucosamine have been
5 disclosed for example, in JP 59013708, WO 98/52576, and U.S. Patent No. 5,866,142, each incorporated herein by reference. To soften and moisturize the skin, a cosmetic containing an N-acetyl amino sugar is disclosed in JP 59013708. A composition for alleviating itching and pain containing N-acetyl-D-glucosamine is disclosed in WO 98/52576. In U.S. Patent No. 5,866,142,
10 a composition for exfoliating the skin has been disclosed, which includes N-acetyl-D-glucosamine. The presence of N-acetyl-D-glucosamine contributes to enhancing the amount of hyaluronic acid which the skin naturally produces in greater quantities in response to exfoliation, induced by other compounds such as histidine. However, no exfoliating activity is attributed to N-acetyl-D-glucosamine. A cosmetic composition containing N-acetylglucosamine and having good adhesion to the skin (i.e., does not lift off of the skin) is disclosed in JP 8188526. However,
15 the prior art does not disclose the ability of glucosamine, and particularly, N-acetyl-D-glucosamine, to exfoliate the skin.

The composition of the present invention contains an exfoliating-effective amount of N-acetyl-D-glucosamine, N-acetylglactosamine, or a combination thereof. By the term "exfoliating-effective amount," as used in the present specification, is meant an amount which
20 effective to cause exfoliation of the skin. The amount of N-acetyl-D-glucosamine, N-acetylglactosamine, or a combination thereof, in the present invention will vary depending on the desired strength or intensity of exfoliation. The N-acetyl-D-glucosamine or N-acetylglactosamine is present in the composition in an amount of about 0.01 to about 25.0 percent of the weight of the composition; preferably 0.5 to about 10.0 percent, and more
25 preferably about 1.0 to 5.0 percent of the weight of the composition.

The N-acetyl-D-glucosamine or N-acetylglactosamine per se is added directly to the cosmetic or pharmaceutical composition by admixing; alternatively, extracts of materials containing substantial quantities of N-acetylglucosamine or N-acetylglactosamine as a component can be used to provide the same concentration. These compositions can also contain
30 chitin. The amounts of chitin in the composition are also present in exfoliating-effective amounts as previously described.

The intensity of exfoliation can also be controlled by the frequency with which the compositions are applied to the skin and the compositions are applied periodically for a period

of time sufficient to exfoliate the skin. Accordingly, the compositions are applied to the skin for a period of at least 2 months, and preferably for at least 4 months, during which time the compositions are applied on a weekly basis. However, a preferred method of obtaining the benefits of the composition is via chronic topical application of the composition to exfoliate the skin. It is suggested, as an example, that "chronic" application be within a range of from about once per week to about 4 to 5 times weekly, preferably daily, most preferably twice daily. By "chronic" application, it is meant herein that the period of topical application may be over the lifetime of the user, preferably for a period of at least about 6 months to about 20 years, more preferably from about 1 year to about 10 years, and still more preferably from about 2 years to about 5 years, thereby resulting in regular desquamation, which may aid in reducing the appearance of fine lines and wrinkles due to chronological aging or photoaging.

The method of the present invention may include applying in addition to the exfoliating effective amount of N-acetyl-D-glucosamine or N-acetylgalactosamine, other optional components, depending on the intended end use of the compositions. These include, but are not limited to, additional exfoliants, preservatives, fragrances, emollients, antiseptics, antiinflammatories, antibacterials, stabilizers, antioxidants, vitamins, pigments, dyes, humectants, and propellants, as well as other classes of materials the presence of which in the compositions may be cosmetically, medicinally, or otherwise desired. Such components can be found in the CTFA International Cosmetics Ingredients Dictionary. Examples of additional exfoliants include but are not limited to chemical exfoliants such as AHAs, for example, lactic acid, or BHAs, for example, salicylic acid, or physical exfoliants such as pumice, polyethylene, walnut shell powder, and the like, or combinations thereof. The amount of additional exfoliants alone or in combination will depend on the type of exfoliant and the strength of exfoliation desired. Preservatives employed, may be in an amount of from about 0.01 to about 2.00 percent, preferably from about 0.02 to about 1.00 percent, of the formula weight. Examples of suitable preservatives are BHA, BHT, propyl paraben, butyl paraben or methyl paraben or an isomer, homolog, analog or derivative thereof.

For topical application, according to the method of the present invention, the compositions can also be formulated with a variety of cosmetically and/or pharmaceutically acceptable vehicles. Accordingly, the compositions of the present invention comprise a pharmaceutically or cosmetically acceptable carrier, in an amount appropriate to accommodate the other components of the formulation. The term "pharmaceutically and/or cosmetically acceptable vehicle" refers to a base, for either pharmaceutical or cosmetic use, within which N-

acetyl-D-glucosamine or N-acetylgalactosamine is soluble and which will not cause harm to humans or other recipients. As used herein, "pharmaceutical" or "cosmetic" will be understood to encompass both human and animal pharmaceuticals or cosmetics. There are few limitations on the type of base which is suitable for the compositions containing N-acetyl-D-glucosamine or N-acetylgalactosamine. The vehicle may be aqueous, nonaqueous or a combination thereof appropriate for the formulation desired.

The compositions can be prepared in any form convenient for topical application to the skin. Such forms include, but are not limited to gels, creams, dispersions, emulsions (water-in-oil or oil-in-water), suspensions, creams, lotions, gels, foams, mousses and the like. In a preferred embodiment, the carrier is a suspension, dispersion or emulsion. The emulsion may be an oil-in-water emulsion, or a water-in-oil emulsion. These emulsions contain one or more oil components, an aqueous component, and a specific emulsifier component chosen depending on the nature of the desired emulsion.

The invention is further illustrated by the following non-limiting examples.

15

EXAMPLE I

A composition, according to the present invention is prepared as follows:

To prepare the composition, the materials are combined in the order above by mixing.

20 The composition is topically applied to the skin for exfoliation.

EXAMPLE II

A composition, according to the present invention is prepared as follows:

25 To prepare the composition, the materials are combined in the order above by mixing. The composition is topically applied to the skin for exfoliation.

EXAMPLE III

A composition according to the present invention, containing N-acetyl-D-glucosamine, is studied to determine the effect on desquamation. A panel of individuals is selected to participate in the test. The selected participants are 15 females between ages 21 and 65 years. Participants are given the composition to take home and self administer on their right hand two times a day, once in the morning after washing and once in the evening at least 15 minutes before

bedtime for 4 weeks. The left hand is the untreated control sample. Participants are allowed to use only the composition being studied and are to log its use in a daily diary. At intervals of 2 and 4 weeks, the participants return for testing without applying the composition for at least 12 hours. The participants are tested without moisturizer or any other product on the skin of their hands.

- 5 Their skin is acclimated to an environment of about 70° F, and about 40 percent relative humidity for about 20 minutes.

- Skin exfoliation is measured with D-Squame discs and image analysis. Specifically, the amount of flakes removed from the skin surface using D-Squame discs are measured. Firmly and evenly, 4 D-Squame discs are pressed on the back of each hand using a hand held uniform
10 pressure device. The discs are removed by gently pulling them away from the skin. After removal, the discs are mounted on clear microscope slides and evaluated with an image analyzer, OPTIMA. A camera takes a picture of the slide and the average gray value is measured to determine the corresponding density of stratum corneocytes. Denser samples have higher gray value differences.
- 15 Results show that the composition reduces skin flakiness after 2 weeks and 4 weeks of treatment when compared to untreated control hand area. After 2 weeks there is about a 15.8 percent decrease and after 4 weeks there was about 16.3 percent decrease in skin flakiness, thereby showing its utility in desquamation.

What we claim is:

1. A method for exfoliation of the skin comprising applying to the skin a composition comprising an exfoliating-effective amount of an amino sugar selected from the group consisting of N-acetyl-D-glucosamine, N-acetylgalactosamine, and a combination thereof.
2. The method according to claim 1 wherein the amino sugar is present in an amount from about 0.01 percent to about 25.0 percent of the weight of the composition.
3. The method according to claim 2 wherein the amino sugar is present in an amount from about 0.5 percent to about 10.0 percent of the weight of the composition.
4. The method according to claim 3 wherein the amino sugar is present in an amount from about 1.0 percent to about 5.0 percent of the weight of the composition.
5. The method according to claim 1 wherein the composition is applied to the skin for a period of at least 4 months.
6. The method according to claim 5 wherein the composition is applied weekly.
7. The method according to claim 1 wherein the composition is applied to the skin for a period of at least 2 months.
8. The method according to claim 1 wherein the compositions are applied from about 4 to about 5 times a week.
9. The method according to claim 8 wherein the composition is applied two times a day.
10. The method according to claim 1 wherein the composition further comprises chitin.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 00/11203

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K7/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 852 946 A (PAOLI AMBROSI GIANFRANCO DE) 15 July 1998 (1998-07-15) abstract page 4, line 37 page 10, line 14 - line 18 -----	1-10
A	WO 97 12597 A (L 'OREAL) 10 April 1997 (1997-04-10) cited in the application the whole document -----	1-10



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z" document member of the same patent family

Date of the actual completion of the international search

28 August 2000

Date of mailing of the international search report

04/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Pelli Wablat, B

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 00/11203

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0852946 A	15-07-1998	IT 85960094 A CA 2219849 A	29-05-1998 29-05-1998
WO 9712597 A	10-04-1997	FR 2739556 A CA 2231553 A EP 0853472 A JP 11501940 T	11-04-1997 10-04-1997 22-07-1998 16-02-1999